Your Exam Manager will receive a copy of the 2014 AIME Solution Pamphlet with the scores.

**CONTACT US --** Correspondence about the problems and solutions for this AIME and orders for any of our publications should be addressed to:

## American Mathematics Competitions 1740 Vine Street Lincoln, NE 68508

Phone: 402-472-2257; Fax: 402-472-6087; email: amcinfo@maa.org

The problems and solutions for this AIME were prepared by the MAA's Committee on the AIME under the direction of:

Jonathan M. Kane AIME Chair kanej@uww.edu


**2014 USA(J)MO --** THE USA MATHEMATICAL OLYMPIAD (USAMO) and the USA MATHEMATICAL JUNIOR OLYMPIAD (USAJMO) are each a 6-question, 9-hour, essay-type examination. The USA(J)MO will be held in your school on Tuesday and Wednesday, April 29 & 30, 2014. Your teacher has more details on who qualifies for the USA(J)MO in the AMC 10/12 and AIME Teachers' Manuals. The best way to prepare for the USA(J)MO is to study previous years of these exams. Copies may be ordered from the web site indicated below.

**PUBLICATIONS** -- For a complete listing of available publications please visit the MAA Bookstore or Competitions site at maa.org.

## The American Mathematics Competitions

are Sponsored by

The Mathematical Association of America - MAA	www.maa.org
The Akamai Foundation	www.akamai.com
Contributors	
Academy of Applied Sciences – AAS	www.aas-world.org
American Institute of Mathematics - AIM	
American Mathematical Association of Two-Year Colleges – AMATYC	www.amatyc.org
American Mathematical Society – AMS	www.ams.org
American Statistical Association – ASA	www.amstat.org
Art of Problem Solving - AoPS	www.artofproblemsolving.com
Association of Symbolic Logic - ASL	
Awesome Math	
Casualty Actuarial Society – CAS	
Conference Board of the Mathematical Sciences - CBMS	www.chmsweh.org
The D.E. Shaw Group	
IDEA Math	
Jane Street Capital	
Math For America	
Math Training Center	
Mu Alpha Theta - MAT	
Pi Mu Epsilon - PME	
Society for Industrial and Applied Math – SIAM	
W.H. Freeman	www.whfreeman.com




- 1. DO NOT OPEN THIS BOOKLET UNTIL YOUR PROCTOR GIVES THE SIGNAL TO BEGIN.
- 2. This is a 15-question, 3-hour examination. All answers are integers ranging from 000 to 999, inclusive. Your score will be the number of correct answers; i.e., there is neither partial credit nor a penalty for wrong answers.
- 3. No aids other than scratch paper, graph paper, ruler, compass, and protractor are permitted. In particular, <u>calculators and computers are not permitted.</u>
- 4. A combination of the AIME and the American Mathematics Contest 12 are used to determine eligibility for participation in the USA Mathematical Olympiad (USAMO). A combination of the AIME and the American Mathematics Contest 10 are used to determine eligibility for participation in the USA Junior Mathematical Olympiad (USAJMO). The USAMO & the USAJMO will be given in your school on TUESDAY and WEDNESDAY, April 29 & 30, 2014.
- 5. Record all of your answers, and certain other information, on the AIME answer form. Only the answer form will be collected from you.

After the contest period, permission to make copies of individual problems in paper or electronic form including posting on web pages for educational use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear the copyright notice.

© 2014 Mathematical Association of America

- 1. Abe can paint the room in 15 hours, Bea can paint 50 percent faster than Abe, and Coe can paint twice as fast as Abe. Abe begins to paint the room and works alone for the first hour and a half. Then Bea joins Abe, and they work together until half the room is painted. Then Coe joins Abe and Bea, and they work together until the entire room is painted. Find the number of minutes after Abe begins for the three of them to finish painting the room.
- 2. Arnold is studying the prevalence of three health risk factors, denoted by A, B, and C, within a population of men. For each of the three factors, the probability that a randomly selected man in the population has only this risk factor (and none of the others) is 0.1. For any two of the three factors, the probability that a randomly selected man has exactly these two risk factors (but not the third) is 0.14. The probability that a randomly selected man has all three risk factors, given that he has A and B is  $\frac{1}{3}$ . The probability that a man has none of the three risk factors given that he does not have risk factor A is  $\frac{p}{q}$ , where p and q are relatively prime positive integers. Find p+q.
- 3. A rectangle has sides of length a and 36. A hinge is installed at each vertex of the rectangle and at the midpoint of each side of length 36. The sides of length a can be pressed toward each other keeping those two sides parallel so the rectangle becomes a convex hexagon as shown. When the figure is a hexagon with the sides of length a parallel and separated by a distance of 24, the hexagon has the same area as the original rectangle. Find  $a^2$ .



4. The repeating decimals  $0.abab\overline{ab}$  and  $0.abcabc\overline{abc}$  satisfy

$$0.abab\overline{ab} + 0.abcabc\overline{abc} = \frac{33}{37},$$

where a, b, and c are (not necessarily distinct) digits. Find the three-digit number abc.

- 13. Ten adults enter a room, remove their shoes, and toss their shoes into a pile. Later, a child randomly pairs each left shoe with a right shoe without regard to which shoes belong together. The probability that for every positive integer k < 5, no collection of k pairs made by the child contains the shoes from exactly k of the adults is  $\frac{m}{n}$ , where m and n are relatively prime positive integers. Find m+n.
- 14. In  $\triangle ABC$ , AB = 10,  $\angle A = 30^{\circ}$ , and  $\angle C = 45^{\circ}$ . Let H, D, and M be points on line  $\overline{BC}$  such that  $\overline{AH} \perp \overline{BC}$ ,  $\angle BAD = \angle CAD$ , and BM = CM. Point N is the midpoint of segment  $\overline{HM}$ , and point P is on ray AD such that  $\overline{PN} \perp \overline{BC}$ . Then  $AP^2 = \frac{m}{n}$ , where m and n are relatively prime positive integers. Find m+n.
- 15. For any integer  $k \geq 1$ , let p(k) be the smallest prime which does not divide k. Define the integer function X(k) to be the product of all primes less than p(k) if p(k) > 2, and X(k) = 1 if p(k) = 2. Let  $\{x_n\}$  be the sequence defined by  $x_0 = 1$ , and  $x_{n+1}X(x_n) = x_np(x_n)$  for  $n \geq 0$ . Find the smallest positive integer, t such that  $x_t = 2090$ .

- 5. Real numbers r and s are roots of  $p(x) = x^3 + ax + b$ , and r + 4 and s 3 are roots of  $q(x) = x^3 + ax + b + 240$ . Find the sum of all possible values of |b|.
- 6. Charles has two six-sided dice. One of the dice is fair, and the other die is biased so that it comes up six with probability  $\frac{2}{3}$ , and each of the other five sides has probability  $\frac{1}{15}$ . Charles chooses one of the two dice at random and rolls it three times. Given that the first two rolls are both sixes, the probability that the third roll will also be a six is  $\frac{p}{q}$ , where p and q are relatively prime positive integers. Find p+q.
- 7. Let  $f(x) = (x^2 + 3x + 2)^{\cos(\pi x)}$ . Find the sum of all positive integers n for which

$$\left| \sum_{k=1}^{n} \log_{10} f(k) \right| = 1.$$

- 8. Circle C with radius 2 has diameter  $\overline{AB}$ . Circle D is internally tangent to circle C at A. Circle E is internally tangent to circle C, externally tangent to circle D, and tangent to  $\overline{AB}$ . The radius of circle D is three times the radius of circle E and can be written in the form  $\sqrt{m}-n$ , where m and n are positive integers. Find m+n.
- 9. Ten chairs are arranged in a circle. Find the number of subsets of this set of chairs that contain at least three adjacent chairs.
- 10. Let z be a complex number with |z|=2014. Let P be the polygon in the complex plane whose vertices are z and every w such that  $\frac{1}{z+w}=\frac{1}{z}+\frac{1}{w}$ . Then the area enclosed by P can be written in the form  $n\sqrt{3}$ , where n is an integer. Find the remainder when n is divided by 1000.
- 11. In  $\triangle RED$ , RD = 1,  $\angle DRE = 75^{\circ}$  and  $\angle RED = 45^{\circ}$ . Let M be the midpoint of segment  $\overline{RD}$ . Point C lies on side  $\overline{ED}$  such that  $\overline{RC} \perp \overline{EM}$ . Extend segment  $\overline{DE}$  through E to point A such that CA = AR. Then  $AE = \frac{a \sqrt{b}}{c}$ , where a and c are relatively prime positive integers, and b is a positive integer. Find a+b+c.
- 12. Suppose that the angles of  $\triangle ABC$  satisfy  $\cos(3A) + \cos(3B) + \cos(3C) = 1$ . Two sides of the triangle have lengths 10 and 13. There is a positive integer m so that the maximum possible length for the remaining side of  $\triangle ABC$  is  $\sqrt{m}$ . Find m.